CENTER FOR DRUG EVALUATION AND RESEARCH APPROVAL PACKAGE FOR:

21-409

APPLICATION NUMBER

Statistical Review(s)

STATISTICAL REVIEW AND EVALUATION CLINICAL STUDIES (STABILITY)

Date NDA# 21-409 Applicant Merck Name of Drug Singulair -Indication Document Reviewed □ Sponsor's cover letter dated 5/21/02: Response to FDA request for information \\CDSESUB1\\N21409\\N 000\\2002-05-21 □ Data sets submitted to EDR: \\CDSESUB1\N21409\N 000\2002-05-21\cmc\datasets Ted J. Guo, Ph.D., Div II/OEB, HFD-715 Statistical Reviewer Prasad Peri, Ph.D., Division of Pulmonary and Allergy Chemist Drug Products (ODE II, HFD-570) Key Words Stability

APPEARS THIS WAY
ON ORIGINAL

Reviewer's Stability Analysis of Singulair

Table 1 describes the batches' characteristics and the upper and lower specifications considered in estimating the expiry-dating period.

Table 1. Description of Singulair and stability analysis

Test parameter	Temperature	Humidity	Strength	Package	Lower spec	Upper spec
Assay	25°C, 30°C	60%	4mg	ī	-	٦
Moisture	25°C, 30°C	60%	4mg	1		1
	25°C, 30°C	60%	4mg	1		1
Total degradation	25°C, 30°C	60%	4mg	1		J

For more information about sponsor's conclusions and CMC reviewer-proposed specification limits, please see Appendix on page 8.

Table 2. Estimate of expiry dating period based on assay (4mg, 25°C, 60%RH)

Fitted Line	Batch	Estimated Expiry Period
Y = 101.0454 + 0.0157 x Time	1005-42	
Y = 100.2666 + 0.0157 x Time	1005-46	
$Y = 99.9635 + 0.0157 \times Time$	MR-4218	
······································	~MIN~	


Figure 1. Estimate of expiry dating period based on assay (4mg, 25°C, 60%RH)

APPEARS THIS WAY ON ORIGINAL

Table 3. Estimate of expiry dating period based on assay (4mg, 30°C, 60%RH)

Fitted Line	Batch	Estimated Expiry Period
Y = 101.1268 + 0.0121 x Time	1005-42	-
Y = 100.5261 + 0.0121 x Time	1005-46	
Y = 100.2894 + 0.0121 x Time	MR-4218	
	~MIN~	

Figure 2. Estimate of expiry dating period based on assay (4mg, 30°C, 60%RH)



APPEARS THIS WAY ON ORIGINAL

Table 4. Estimate of expiry dating period based on moisture (4mg, 25°C, 60%RH)

Fitted Line	Batch	Estimated Expiry Period
$Y = 0.1931 + 0.0002 \times Time$	POOLED	
	-MIN-	

Figure 3. Estimate of expiry dating period based on moisture (4mg, 25°C, 60%RH)




Table 5. Estimate of expiry dating period based on moisture (4mg, 30°C, 60%RH)

Fitted Line	Batch	Estimated Expiry Period
$Y = 0.1817 + 0.0004 \times Time$	POOLED	
	-MIN-	

Figure 4. Estimate of expiry dating period based on moisture (4mg, 30°C, 60%RH)

APPEARS THIS WAY ON ORIGINAL

itted Line	Batch	Estimated Expiry Period	
= 0.3239 + 0.0004 x Time	POOLED	· · · · · · · · · · · · · · · · · · ·	
	~MIN~		
igure 5. Estimate of expiry	dating period ba	sed on — (4mg, 25°C, 60%)	UH)
		7	
)	
		}	
able 7. Estimate of expiry (dating period base	ل d on —— (4mg, 30°C, 60%R)	(F
itted Line	Batch	Estimated Expiry Period	•
= 0.3591 + 0.0010 x Time	1005-42		
= 0.2308 + 0.0010 x Time	1005-46		
= 0.3083 + 0.0010 x Time	MR-4218		
	~MINI~		
	~MIN~	4 - 2005 (00)	N. F. P. N.
igure 6. Estimate of expiry		sed on (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on (4mg, 30°C, 60%1	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)
igure 6. Estimate of expiry		sed on — (4mg, 30°C, 60%)	RH)

Table 8. Estimate of expiry dating period based on total degradation (4mg, 25°C, 60%RH)

Fitted Line	Batch	Estimated Expiry Period
Y = 0.3323 + 0.0002 x Time	POOLED	
***************************************	-MIN-	

Figure 7. Estimate of expiry dating period based on total degradation (4mg, 25°C, 60%RH)

Table 9. Estimate of expiry dating period based on total degradation (4mg, 30°C, 60%RH)

Fitted Line	Batch	Estimated Expiry Period
Y = 0.3391 + 0.0018 x Time	1005-42	-
Y = 0.3073 - 0.0007 x Time	1005-46	
Y = 0.3079 + 0.0011 x Time	MR-4218	,
	~MIN~	

Figure 8. Estimate of expiry dating period based on total degradation (4mg, 30°C, 60%RH)

APPEARS THIS WAY ON ORIGINAL

Table 10. Estimated expiry-dating period

Test parameter	Condition	Estimated expiry-dating period (in weeks)
Assay	25°C, 60%RH	
Assay	30°C, 60%RH	
Moisture	25°C, 60%RH	
Moisture	30°C, 60%RH	
	25°C, 60%RH	
	30°C, 60%RH	تب
degradation	25°C, 60%RH	
degradation	30°C, 60%RH	
	~MIN~	<u>, , , , , , , , , , , , , , , , , , , </u>

Conclusion

The data show that the estimated expiry-dating period, with 95% confidence, is at least — weeks. The sponsor proposed expiry dating period was supported by the data.

APPEARS THIS WAY

Redacted _____

page(s) of trade secret

and/or confidential

commercial information

(b4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

Ted Guo 7/1/02 01:31:44 PM BIOMETRICS

Lisa A. Kammerman 7/1/02 03:22:06 PM BIOMETRICS I concur with review

APPEARS THIS WAY ON ORIGINAL